

THE EFFECT OF VIBRATION ON THE MORPHOLOGIC
PATTERN OF ENDOCRINE GLANDS

Ryszard Piechocinski

NASA TT F-11,328

FACILITY FORM 602

N67-40465

(ACCESSION NUMBER)

3 (PAGES)

(THRU)

1 (CODE)

(NASA CR OR TMX OR AD NUMBER)

(CATEGORY) 04

Translation of "Wplyw Wibracji Na Morfologiczny Orbaz Gruczolów Dokrewnych."
Patologia Polska, Vol. 17, No. 4, pp 561-563, 1966

NATIONAL AERONAUTICS AND SPACE ADMINISTRATION
WASHINGTON, D. C. 20546 OCTOBER 1967

THE EFFECT OF VIBRATION ON THE MORPHOLOGIC
PATTERN OF ENDOCRINE GLANDS

Ryszard Piechocinski

Received from the Institute of General and Experimental
Pathology of the Medical Academy in Cracow
Director: Prof. B. Giedosz, M.D.

ABSTRACT. After vibration of frequency 1800/min and amplitude of 6 mm in the vertical and 4 mm in the horizontal direction, no changes were found in the thyroid gland or ovaries. The adrenal glands were enlarged, their medullary portion was hyperemic, and the zona fasciculata was slightly enlarged. The changes in the adrenal glands of white mice were attributed to stress connected with vibration.

As found in numerous reports, various changes appear in the organism under the influence of vibration. Changes were observed in joints, bones, muscles, vessels etc. Intensity and localization of these changes, as well as the time at which they appear, depend on the frequency, amplitude, and duration of vibration. Recently, biochemic changes due to vibration have been noticed. Thus, for example, Stanosek et al. in an experiment with animals discovered changes in the activity of alanine and asparagin aminotransferase at 5 hertz and an amplitude of 20 mm. Stoklosa demonstrated changes in the behavior of phosphatases in the parenchymal organs of animals. Babadzianian et al. described disorders in the menstrual period in women who worked with compressed-air hammers. Komorowska and Krolikowska analyzed the relation between the frequency of miscarriages and premature childbirths, and the kind of work performed by Lodz** female textile workers. Vibrations, therefore, can cause various changes in different tissues and organs. /561*

In the literature accessible to us, we have not found any information on the effect of vibrations on endocrine glands. In the experiment presented below, we therefore decided to find out whether vibrations can cause changes in these glands. We have studied only some endocrine glands.

Materials and Methods

In the experiment we have used white virgin mice, 20-22g in weight, of the same breed, identically fed and maintained under the same environmental conditions. We utilized vibration devices of 1800 oscillations per second, with an amplitude of 6 mm in the vertical direction and 4 mm in the horizontal direction. The first group consisted of control mice. The second group, composed of 30 mice, was subjected to vibrations twice a day for 15 minutes for 10 days. The third group, 30 mice, was subjected to vibrations three times a day for 30 days, 10 minutes each day. Samples were taken of the thyroid gland,

* Note: Numbers in the margin indicate pagination in the original foreign text.

** Translator's Note: Lodz is Poland's main center of the textile industry.

ovaries, and the adrenal glands. The thyroid gland was fixed in alcohol-formol and paraffin strips were dyed with hematoxylin and eosin. Ovaries were fixed in 10% formol. Paraffin strips were made from one ovary, and subsequently dyed with hematoxylin and eosin. From the other ovary, frozen strips were dyed with scarlet R according to the method by Hexheimer. The adrenal glands were treated in a similar way.

Results of the Investigations

The thyroid glands of mice from both groups subjected to vibrations have not shown any histological changes, as compared with the pattern for thyroid glands of the control animals.

No deviations from the norm were found in survey specimens prepared from the ovaries of both groups of animals subjected to vibrations. Similarly, no differences were detected in the amount and distribution of lipids dyed with scarlet R.

The adrenal glands of mice subjected to vibrations were macroscopically larger than those of the mice from the control group. In the survey specimens made from the adrenal glands of animals subjected to vibrations, we found hyperemia of the medullary portion of the adrenal glands, and sometimes blood on the outside of the vessels. Furthermore, we discovered that the zona fasciculata of the adrenal glands was slightly enlarged. No differences were detected in the amount and distribution of lipids.

Analysis of the Results

Under the action of vibration described above, we have not found any changes in the thyroid gland and ovaries. The adrenal glands were slightly enlarged, the medullary portion was hyperemic, and the zona fasciculata was somewhat wider than normal. These changes are considered to be a result of the adaptation reaction, with vibrations being a serious stress to the organism.

REFERENCES

1. Babadzanian, M. G., Kostina, E. I., Chirkov, W. J. A Study of Working Conditions and Physiological Disturbances in Female Workers Employed as Crosstie Movers, *Gigiena i Sanitaria*, 2, 21, 1957.
2. Bugyi, B. Bone Changes in Blacksmiths. Contribution to the Illnesses of Compressed Air Device Operators, *Zentralblatt für Arbeitsmedizin und Arbeitsschutz*, 2, 1957.
3. Komorowska, A., Krolikowska, M. The Effect of Work on the Frequency and Course of Miscarriages and Premature Childbirths, *Medycyna Pracy*, 1, 217, 1961.
4. Stanosek, J., Gregorczyk, J., Lewandowska-Tokarz, A. Studies on the Effect of Prolonged Vertical Vibrations of a Low Frequency on Biochemical Disturbances in Guinea Pigs, *Med. Dosw. i Mikr.*, 4, 347, 1964.
5. Stoklosa, E., Ludyga, K. The Effect of Mechanical Vibrations of Lower Frequencies on the Weight of Growing Rats, *Post. Hig. i Med. Dosw.*, 17, 781, 1963.

/563

6. Stoklosa, E. Histochemical Changes in Parenchymal Organs of White Mice, Subjected to the Influence of Mechanical Vibrations of Low Frequencies, Post. Hig. i. Med. Dosw., 17, 785, 793, 799, 1963.

Scientific Translation Service
4849 Tocaloma Lane
La Canada, California 91011

NASw 1496